

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
29 April 2004 (29.04.2004)

PCT

(10) International Publication Number
WO 2004/035026 A1

(51) International Patent Classification⁷: **A61K 9/20**,
31/41, A61P 39/04

CASSIERE, Jean-Pierre [FR/FR]; 58-60 route de
l'Empereur, F-92500 Rueil Malmaison (FR).

(21) International Application Number:
PCT/EP2003/011351

(74) Agent: **GRUBB, Philip**; Novartis AG, Corporate Intellectual
Property, CH-4002 Basel (CH).

(22) International Filing Date: 14 October 2003 (14.10.2003)

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE,
GH, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,
LK, LT, LU, LV, MA, MD, MK, MN, MX, NI, NO, NZ,
OM, PG, PH, PL, PT, RO, RU, SC, SE, SG, SK, SY, TJ,
TM, TN, TR, TT, UA, US, UZ, VC, VN, YU, ZA, ZW.

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
0223978.8 15 October 2002 (15.10.2002) GB

(84) Designated States (*regional*): Eurasian patent (AM, AZ,
BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE,
BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU,
IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR).

(71) Applicant (*for all designated States except AT, US*): **NO-
VARTIS AG** [CH/CH]; Lichtstrasse 35, CH-4056 Basel
(CH).

Published:

- with international search report
- before the expiration of the time limit for amending the
claims and to be republished in the event of receipt of
amendments

(71) Applicant (*for AT only*): **NOVARTIS PHARMA GMBH**
[AT/AT]; Brunner Strasse 59, A-1230 Vienna (AT).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **DEFPEZ, Karine**
[FR/FR]; 57, rue du Rhône, F-69007 Lyon (FR).

*For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.*

(54) Title: DEFERACIROX DISPERSIBLE TABLETS

(57) Abstract: The invention pertains to dispersible tablets comprising as active ingredient 4-[3,5-bis(2-hydroxyphenyl)-[1,2,4] triazol-1-yl] benzoic acid or pharmaceutically acceptable salt thereof in an amount of from 5 to 40% in weight by weight of the total tablet.



WO 2004/035026 A1

DEFERASIROX DISPERSIBLE TABLETS

The present invention relates to dispersible tablets, e.g. pharmaceutical dispersible tablets, comprising 4-[3,5-bis(2-hydroxyphenyl)-[1,2,4]triazol-1-yl]benzoic acid or a pharmaceutically acceptable salt thereof, and is hereinafter referred to as Compound I.


Compound I is an orally active iron chelator that is indicated in the treatment of iron overload in transfusion dependent anemias, in particular thalassemia major, thalassemia intermediate and in sickle cell disease to reduce iron-related morbidity and mortality. Compound I can also be used in the treatment of hemochromatosis.

Clinical thalassemia (major and intermedia) are hereditary disorders characterized by defective production of hemoglobin, which leads to decreased production and increased destruction of red blood cells.

Sickle cell disease is caused by a mutation in the hemoglobin-Beta gene leading to the production of abnormal hemoglobin S. Normal red blood cells die after 120 days and sickle cells (red blood cells with hemoglobin S) are destroyed more rapidly (10 to 20 days) causing anemia. This anemia is what gives the disease its commonly known name - sickle cell anemia.

Hemochromatosis, the most common form of iron overload disease, is an inherited disorder that causes the body to absorb and store too much iron. The extra iron builds up in organs and damages them. Without treatment, the disease can cause these organs to fail.

Patients with sickle cell disease or thalassemia, who receive significant numbers of blood transfusions and patients with hemochromatosis require therapy to remove iron from the body, called chelation therapy.



Compound I has the following formula:

- 2 -

Compound I in the free acid form, salts thereof and its crystalline forms are disclosed in the International Patent Publication WO 97/49395, which is hereby incorporated by reference (published on December 31, 1997).

Typically, prescribed daily dosages of Compound I for the treatment of thalassemia are high, e.g. 5 to 40 mg/kg of body weight/day in adults or children. In children, the dosage is preferably 5 to 30 mg/kg of body weight/day. Due to the high dosage strength, the tablet dimensions do not permit the formulation of a conventional tablet. Thus, there is a need for an oral dosage form that is convenient to administer to adults and to children and that provides a pharmacologically active daily dosage amount of Compound I.

Present inventors have now surprisingly found that the formulation of Compound I in form of a dispersible tablet allows an oral dosage form with a high drug loading and which is convenient to administer to, for example children and elderly, and stable.

By "dispersible tablet" is meant a tablet which disperses in aqueous phase, e.g. in water, before administration.

Accordingly, the present invention provides a dispersible tablet with high drug loading comprising Compound I as active ingredient, the active ingredient being present in an amount of from about 5% to 40%, e.g. at least about 10, 15, 20 or 25 %, preferably more than 25% in weight based on the total weight of the dispersible tablet. In particular, the amount of Compound I may vary from 25 to 40%, e.g. 28 to 32% in weight based on the total weight of the dispersible tablet.

The present invention pertains to a dispersible tablet comprising an iron-chelating pharmacologically effective amount of Compound I or a pharmaceutically acceptable salt thereof present in an amount of from 5% to 40% weight by weight based on the total weight of the tablet.

In one aspect of the invention is provided a dispersible tablet comprising Compound I or a pharmaceutically acceptable salt thereof present in an amount of from 5% to 40% in weight based on the total weight of the tablet.

Compound I may be in the free acid form or pharmaceutically acceptable salts thereof, preferably in the free acid form. The active moiety corresponds to Compound I in the free acid form. Within the context of this disclosure, reference to Compound I is understood to include Compound I in its free acid form or a pharmaceutically acceptable salt thereof or any crystal forms thereof including hydrates or solvates, if not indicated otherwise and where appropriate and expedient.

The present invention also provides a dispersible tablet comprising:

- (a) Compound I or a pharmaceutically acceptable salt thereof, and
- (b) at least one pharmaceutically acceptable excipient suitable for the preparation of dispersible tablets wherein the amount of Compound I or a pharmaceutically acceptable salt thereof, calculated as the percentage of the content in weight of the active moiety based on the total weight of the dispersible tablet, is from about 5% to 40% t, e.g. at least about 10, 15, 20 or 25 %, preferably more than 25% in weight based on the total weight of the dispersible table. In particular, the amount of Compound I as active ingredient may vary from 25 to 40%, e.g. 28 to 32% in weight based on the total weight of the dispersible tablet.

In a preferred embodiment of the invention, the present invention provides a dispersible tablet wherein Compound I is in the free acid form (Compound I free acid form).

In a most preferred aspect of the invention, Compound I in the free acid form is in a crystalline form.

One or more pharmaceutically acceptable excipients may be present in the dispersible tablets, e.g. those conventionally used, e.g. (1.1) at least one filler, e.g., lactose, ethylcellulose, microcrystalline cellulose, (1.2) at least one disintegrant, e.g. cross-linked polyvinylpyrrolidinone, e.g. Crospovidone®, (1.3) at least one binder, e.g. polyvinylpyrrolidinone, hydroxypropylmethyl cellulose, (1.4) at least one surfactant, e.g. sodium laurylsulfate, (1.5) at least one glidant, e.g. colloidal silicon dioxide, (1.6), at least one lubricant, e.g. magnesium stearate.

Reference is made to the extensive literature on the subject for these and other pharmaceutically acceptable excipients and procedures mentioned herein, see in particular Handbook of Pharmaceutical Excipients, Third Edition, edited by Arthur H. Kibbe, American

Pharmaceutical Association, Washington, USA and Pharmaceutical Press, London; and Lexikon der Hilfsstoffe für Pharmazie, Kosmetik and angrenzende Gebiete edited by H.P. Fiedler, 4th Edition, Editio Cantor, Aulendorf and earlier editions which are incorporated herein by reference.

Fillers (1.1) according to the invention are lactose especially lactose monohydrate, preferably lactose monohydrate (200mesh) and lactose spray dried, microcrystalline cellulose especially PH 102, PH 101.

Suitable disintegrants (1.2) according to the invention include but are not restricted to maize starch, CMC-Ca, CMC-Na, microcrystalline cellulose, cross-linked PVP, e.g. as known and commercially available under the trade names Crospovidone®, Polyplasdone®, available commercially from the ISP company, or Kollidon® XL, alginic acid, sodium alginate and guar gum. Preferably, cross-linked PVP, e.g. Crospovidone® is used.

Binders (1.3) include but are not restricted to starches, e.g. potato, wheat or corn starch, microcrystalline cellulose, e.g. products such as Avicel®, Filtrak®, Heweten® or Pharmacel®; hydroxypropyl cellulose, hydroxyethyl cellulose, hydroxypropylmethyl cellulose, e.g. hydroxypropylmethyl cellulose-Type 2910 USP, hypromellose, and polyvinylpyrrolidone, e.g. Povidone® K30 from BASF. Preferably, polyvinylpyrrolidone is used, most preferably PVP K.30.

Appropriate surfactant (1.4) according to the invention may be used: sodium laurylsulfate, betain, quaternary ammonium salts, polysorbates, sorbitan esters and poloxamer. Preferably the surfactant is sodium laurylsulfate.

As glidants (1.5), one or more of the following may be used: silica; colloidal silica, e.g. colloidal silica anhydrous, e.g. Aerosil® 200, magnesium trisilicate, powdered cellulose, starch and talc. Preferably, colloidal silicon dioxide is used.

As lubricants (1.6) one or more of the following may be used Mg-, Al- or Ca-stearate, PEG 4000 – 8000, talc, sodium benzoate, glyceryl mono fatty acid, e.g. having a molecular weight of from 200 to 800 Daltons, e.g. glyceryl monostearate (e.g. Danisco, UK), glyceryl dibehenate (e.g. CompritolATO888™, Gattefossé France), glyceryl palmito-stearic ester (e.g. Precirol™,

Gattefossé France), polyoxyethylene glycol (PEG, BASF), hydrogenated cotton seed oil (Lubitrab, Edward Mendell Co Inc), castor seed oil (Cutina HR, Henkel). Preferably, magnesium stearate is used.

One or more of these pharmaceutically acceptable excipients may be selected and used having regard to the particular desired properties of the dispersible tablet by routine experimentation.

According to the present invention, the amount of filler (1.1) may vary within a range of from about 35 to 55%, in particular 40 to 50% in weight based on the total weight of the dispersible tablet.

The amount of disintegrant (1.2) may vary within a range of from 5 to 40%, e.g. 10 to 35% in weight based on the total weight of the dispersible tablet.

The amount of binder (1.3) may vary from 1 to 10 %, preferably from 1.5 to 5 % in weight based on the total weight of the dispersible tablet.

The amount of surfactant (1.4) may vary from 0.1 to 2%, preferably from 0.2 to 1%.

The amount of glidant (1.5) may vary within ranges of from 0.1 to 5%, in particular 0.1 to 2.5%, e.g. 0.1 to 0.5% in weight based on the total weight of the dispersible tablet.

The amount of lubricant (1.6) may be below 1% in weight based on the total weight of the dispersible tablet, preferably below 0.5%, most preferably below 0.4% and even most preferably the amount of lubricant is ranging between 0.01% and 0.4%. Very preferably the amount of lubricant is above 0.02% and below 0.4% in weight based on the total weight of the dispersible tablet.

It will be appreciated that any given excipient may serve more than one function e.g. as filler, disintegrant, binder, glidant, and/or lubricant.

In one aspect of the invention, a lubricant is present in less than 1% in weight based on the total weight of the dispersible tablet, preferably less than 0.4%.

The invention also pertains to a dispersible tablet wherein the lubricant is magnesium stearate.

In a preferred aspect of the invention, the dispersible tablet comprises the following pharmaceutically acceptable excipients: one or more fillers in a total amount of about 40% to 50% in weight based on the total weight of the dispersible tablet, one or more binders in a

total amount of about 1.5% to 5% in weight based on the total weight of the dispersible tablet, one or more disintegrants in a total amount of about 10% to 35% in weight based on the total weight of the dispersible tablet, one or more glidants in a total amount of about 0.1% to 0.5% in weight based on the total weight of the dispersible tablet, and/or one or more lubricants in a total amount of about 0.01 % to 0.4 % in weight based on the total weight of the dispersible tablet.

In a preferred aspect of the invention, the dispersible tablet comprises the following pharmaceutically acceptable excipients: one or more fillers in a total amount of about 40% to 50% in weight based on the total weight of the dispersible tablet, one or more binders in a total amount of about 1.5% to 5% in weight based on the total weight of the dispersible tablet, one or more disintegrants in a total amount of about 10% to 35% in weight based on the total weight of the dispersible tablet, one or more surfactant in a total amount of about 0,2% to 1% in weight based on the total weight of the dispersible tablet, one or more glidants in a total amount of about 0.1% to 0.5% in weight based on the total weight of the dispersible tablet, and/or one or more lubricants in a total amount of about 0.01 % to 0.4 % in weight based on the total weight of the dispersible tablet.

The absolute amounts of each pharmaceutically acceptable excipient and the amounts relative to other pharmaceutically acceptable excipients is similarly dependent on the desired properties of the dispersible tablet and may also be chosen by routine experimentation.

The present inventors have encountered difficulties in the production of dispersible tablets comprising Compound I which may be due to the low density of the active ingredient, to its electrostatic characteristics which may lead to a poor flowability and to its sticking tendency.

In accordance with the present invention, it has now unexpectedly been found that pharmaceutically acceptable oral solid dosage forms in the forms of dispersible tablets convenient for patient administration and dispersible in 5 minutes or less, preferably 3 minutes or less, may be obtained by preparation of tablets by compression methods. More specifically, the dispersible tablets of the invention may be prepared by granulation, preferably wet-granulation, followed by compression methods, preferably under spray lubrication.

In general, wet-granulation may be used to improve flowability and sticking tendency, however, wet-granulation process is not preferred when the pharmaceutical composition is to be a dispersible tablet. Indeed, wet-granulation increases the cohesion of the active ingredient particles and increases the disintegration time of the final tablet which is not in accordance with patient compliance or the European Pharmacopoeia which requests a disintegration time of 3 minutes or less for a dispersible tablet. Moreover, present inventors have encountered the problem that even upon wet-granulation the active ingredient remains sticky and is difficult to handle in a tableting machine. Present inventors have now surprisingly found that stickiness may be resolved, e.g. by adding lubricant to the composition of the tablet without increasing the disintegration time, e.g. above acceptable values, e.g. above 5 minutes.

The dispersible tablets of the invention have a disintegration time, e.g. in aqueous media, e.g. in water, of 5 minutes or below 5 minutes. The dispersible tablets of the invention are, despite the high drug loading, dispersible, e.g. in aqueous media, e.g. in water, in less than 5 minutes, preferably less than 3 minutes, and, therefore, convenient to administer, e.g. to children or elderly. This leads to a better patient compliance.

In another embodiment, this invention provides a dispersible tablet comprising from 100 mg to 800 mg of Compound I as active ingredient, e.g. of from 100 mg to about 600 mg. Most preferably, dispersible tablets according to the invention are dispersible tablets containing 125 mg, 250 mg or 500 mg of Compound I as active ingredient.

Accordingly, the present invention provides dispersible tablets, e.g. dispersible tablets, containing an amount of Compound I, equal to 125 mg, 250 mg, or 500 mg of Compound I free acid form. Most preferably, the Compound I in the free acid form used for the dispersible tablet according to the invention is the crystalline form, especially the crystalline form the preparation of which is described in example 5 of WO 97/49395, which is hereby incorporated by reference.

According to the invention, the process for the preparation of the dispersible tablets consists of granulating an inner phase, mixing (together) it with one or more pharmaceutically acceptable excipient(s) and compressing the obtained mixture under spray lubrication conditions.

The inner phase comprises Compound I. Preferably, the inner phase comprises Compound I and one or more pharmaceutically acceptable excipients. Preferably, the pharmaceutically acceptable excipients of the inner phase are one or more fillers, one or more disintegrants, one or more binders and one or more surfactants. Preferably the amount of one or more fillers in the inner phase is ranging from about 5 to 35% in weight based on the total weight of the dispersible tablet, more preferably 10 to 30% and most preferably 15 to 25%. The filler according to the invention is preferably lactose monohydrate. The disintegrant is preferably Croscopovidone XL. The amount of disintegrant present in the inner phase is preferably ranging from 5 to 30%, more preferably 7 to 25% in weight based on the total weight of the dispersible tablet. The Compound I and one or more fillers and one or more disintegrants are mixed together with a wetting solution comprising one or more surfactants, water and one or more binders. The preferred binder is PVP K.30. The mixture is processed for granulation, e.g. using a wet high-shear granulator to form the wet-granulates. The wet-granulates may then be dried, e.g. using a fluid bed dryer, and calibrated, e.g. using an oscillating granulator. The outer phase consists of one or more pharmaceutically acceptable excipient(s) and is mixed with the inner phase using e.g. a free fall mixer. Preferably, one or more fillers and one or more glidants are added. Most preferably, cellulose microcrystalline and lactose are added as fillers. Even more preferably, microcrystalline cellulose is added in the range of 5 to 20% in weight based on the total weight of the dispersible tablet and lactose is added in the range of 5 to 20% in weight based on the total weight of the dispersible tablet. The outer phase according to the invention may also contain one or more glidants, most preferably colloidal silicon dioxide. In a preferred embodiment, the amount of glidant in the outer phase is ranging from about 0.1 to 5%, preferably 0.1 to 2.5%, most preferably 0.1 to 0.5% in weight based on the total weight of the tablet.

In one aspect of the invention one or more lubricants, instead of being incorporated into the mixture of the inner and outer phase, may be deposited on the punches of the tableting machine before compression. According to the invention, one or more lubricants may be sprayed on the material contacting surfaces of pressing tools, e.g. punches and/or dies, of the tableting machine before compression. Preferably, one or more lubricants are sprayed on the material contacting surfaces of pressing tools, e.g. punches and dies, of the tableting machine before compression.

In one embodiment of the invention, the process for the preparation of a dispersible tablet comprises

- (a) forming an inner phase comprising
 - (i) mixing Compound I together with pharmaceutically acceptable pharmaceutically acceptable excipients,
 - (ii) wet-granulating the mixture obtained in (i);
- (b) forming an outer phase comprising
 - (iii) adding further pharmaceutically acceptable excipients to the inner phase obtained in (ii) and mixing;
- (c) forming the dispersible tablet by
 - (iv) compressing the mixture obtained in step (iii) under spray lubrication condition.

In a further aspect the present invention provides a process comprising:

- (i) mixing Compound I and pharmaceutically acceptable excipients, e.g. one or more fillers, e.g. lactose, and one or more disintegrants, e.g. Crospovidone XL, in a high shear mixer;
- (ii) adding a solution of one or more surfactant and one or more binder, subjecting the mixture to wetting/kneading, e.g. in a high shear mixer, wet-granulating using, e.g. a rotating impeller, drying, e.g. in a fluidized bed dryer, then calibrating in an oscillating granulator, and;
- (iii) adding pharmaceutically acceptable excipients, e.g. sieved excipients, such as one or more fillers, e.g. microcrystalline cellulose or lactose, one or more glidant, e.g. colloidal silicon dioxide, and mixing, e.g. in a free fall mixer;
- (iv) tableting the mixture obtained in step (iii) by compression, e.g. in a conventional tablet press, preferably a rotary machine and spraying the lubricant on the materials contacting surfaces of pressing tools.

Procedures which may be used may be conventional or known in the art or based on such procedures e.g. those described in L. Lachman et al. *The Theory and Practice of Industrial Pharmacy*, 3rd Ed, 1986, H. Sucker et al, *Pharmazeutische Technologie*, Thieme, 1991, Hagers *Handbuch der pharmazeutischen Praxis*, 4th Ed. (Springer Verlag, 1971) and Remington's *Pharmaceutical Sciences*, 13th Ed., (Mack Publ., Co., 1970) or later editions.

By "inner phase" is meant the granulate phase (steps (i) and (ii)) including the active ingredient Compound I and one or more the pharmaceutically acceptable excipients.

By "outer phase" is meant one or more pharmaceutically acceptable excipients added to the inner phase (granulates) (step (iii)).

By "total weight of the dispersible tablet" is meant the weight of a tablet being the inner and the outer phase.

The physical and chemical stability may be tested in conventional manner, e.g. the dispersible tablets may be tested as such by measurement of dissolution, friability, disintegration time, assay for Compound I degradation products, appearance and/or microscopy, e.g. after storage at room temperature, i.e. 25°C, and/or storage at 40°C.

The dispersible tablets may vary in shape and be, for example, round, oval, oblong, cylindrical or any other suitable shape. In one aspect, the dispersible tablets according to the invention contain small amount of magnesium stearate, e.g. about 0.01 % to 0.4 % in weight based on the total weight of the dispersible tablet, having regard to the amount of Compound I contained therein, thus allowing a disintegration time, which complies with the European Pharmacopoeia Specifications.

In a preferred embodiment of the invention dispersible tablets obtained by the compression method described above are round or oval. The edges of the dispersible tablets may be beveled or rounded. Most preferably, the dispersible tablets are round with bevelled edges. The dispersible tablets according to the invention may be scored, embossed or engraved.

The dispersible tablet according to the invention is preferably round, flat with bevelled edges. The 125 mg dispersible tablet has a diameter ranging between 10 and 20 mm, most preferably between 10 and 15 mm. The preferred diameter of the 125 mg dispersible tablet is 12 mm. Its thickness is ranging from 2.5 to 4.5 mm, preferably between 3.2 and 3.9 mm. The 250 mg dispersible tablet has a diameter ranging from 12 to 20 mm, preferably between 14 and 18 mm, the most preferred diameter is 15 mm. Its thickness is ranging from 3.5 to 5.5 mm, most preferably between 4 and 5 mm. The 500 mg dispersible tablet has a diameter ranging from

15 to 30 mm, preferably between 15 and 25 mm, the most preferred diameter is 20 mm. Its thickness is ranging from 4.5 to 6.5 mm, most preferably between 5 and 6 mm.

The dispersible tablets of the invention comprising about 125 mg of Compound I as active moiety may have a hardness of from about 50 to 120 N, preferably 60 to 100 N.

The dispersible tablets of the invention comprising about 250 mg of Compound I may have a hardness of 70 to 150 N, preferably 90 to 130 N. The dispersible tablets of the invention comprising about 500 mg of Compound I may have a hardness of 80 to 190 N, preferably 110 to 160 N.

Preferably, the disintegration time is not more than 5 minutes, most preferably the disintegration time is less than 3 minutes as measured using a disintegration time apparatus.

By "disintegration time" is meant the time that needs the dispersible tablet to disintegrate in water at room temperature in a disintegration time device.

The dispersible tablet of the present invention is dispersible in an aqueous phase, preferably water.

The dispersible tablets of the invention may be colored and/or marked so as to impart an individual appearance and to make them instantly recognizable. The use of dyes can serve to enhance the appearance as well as to identify the dispersible tablets. Dyes suitable for use in pharmacy typically include carotinoids, iron oxides or chlorophyll. The dispersible tablets of the invention may be marked using an imprint code.

The dispersible tablets of the invention are useful for the treatment of iron overload in transfusion dependent anemias, in particular thalassemia major, thalassemia intermediate and sickle cell disease and in the treatment of hemochromatosis.

The activity and characteristics of the dispersible tablets of the invention may be indicated in standard clinical trials and/or animal trials.

The dispersible tablets of the invention are stable both to the production process and during storage, e.g. for 2 years or even 3 years in conventional packaging, e.g. sealed aluminium

blister packs or triplex blister packs. Less than about 5%, e.g. 2 or 3% or less of Compound I as active ingredient may degrade during this time as determined in conventional tests. For example, less than 1% of Compound I as active ingredient is degraded in one year in HDPE filled bottles.

Depending on age, individual condition, mode of administration, and the clinical picture in question, effective daily dosing, e.g. 350 to 2800 mg of Compound I, are administered to patients of 70 kg body weight.

The invention further relates also to a method of administering to a mammal, preferably a human subject, in need of such a treatment, Compound I in the form of a dispersible tablet. The invention relates especially to such method wherein a daily dose of 5 to 40 mg/kg of body weight of Compound I as active ingredient is administered to a patient. It will be understood that the specific dose level for any particular patient will depend upon a variety of factors including the age, the body weight, general health, drug combination with one or more active drugs, type and severity of the disease.

The invention further provides a medicament package comprising dispersible tablets according to the invention and printed instructions directing that one or more dispersible tablets of Compound I be administered orally.

The following non-limitative examples illustrate the invention.

- 13 -

Example 1: Dispersible tablet Formulation (125 mg, 250 mg and 500 mg dispersible tablets)
with a disintegration time above 3 minutes.

		Amount per dispersible tablet (mg)			
Components		%	125 mg	250 mg	500 mg
Phase I	Compound I (free acid form)	29.41	125.0	250.00	500.0
	Lactose 200 Mesh (1.1)	22.09	93.88	187.75	375.5
	Crospovidone XL (1.2)	10.00	42.50	85.00	170.00
Phase II	PVP K.30 (1.3)	3.00	12.75	25.50	51.00
	Sodium laurylsulfate (1.4)	0.50	2.13	4.25	8.50
Phase III	Crospovidone XL (1.2)	10.00	42.50	85.00	170.00
	Microcrystalline cellulose (1.1)	11.90	50.57	101.15	202.3
	Lactose spray dried (1.1)	11.90	50.57	101.15	202.3
	Aerosil 200 (1.5)	0.20	0.85	1.70	3.40
Phase IV	Magnesium stearate (1.6)	1.00	4.25	8.50	17.00
Tablet weight (mg)		100.00	425.00	850.00	1700.00
Tablet diameter (mm)		-	12	15	20

Example 2: Dispersible tablet Formulation (125 mg, 250 mg and 500 mg dispersible tablets) with a disintegration time below 3 minutes.

Components		%	Amount per dispersible tablet (mg)		
			125 mg	250 mg	500 mg
Phase I	Compound I (free acid form)	29.4	125.0	250.0	500.0
	Lactose 200 Mesh (1.1)	17.1	72.6	145.2	290.4
	Crospovidone XL (1.2)	15.0	63.7	127.4	254.8
Phase II	PVP K.30 (1.3)	3.0	12.8	25.6	51.2
	Sodium laurylsulfate (1.4)	0.5	2.1	4.2	8.4
Phase III	Crospovidone XL (1.2)	5.0	21.3	42.6	85.2
	Microcrystalline cellulose (1.1)	14.9	63.3	126.6	253.2
	Lactose spray dried (1.1)	14.9	63.3	126.6	253.2
	Aerosil 200 (1.5)	0.2	0.9	1.8	3.6
Phase IV	Magnesium stearate (1.6)	< 0.2*			
Tablet weight (mg)		100.0	425	850	1700
Tablet diameter (mm)		-	12	15	20
Tablet thickness (mm)		-	3.6+/-0.3	4.5+/-0.3	5.5+/-0.3

Dispersible tablets of Compound I free acid according to the invention are prepared by forming a inner phase by wet granulation of a mixture of Phase I and Phase II ingredients, Phase III ingredients formed the outer phase and the lubricant (Phase IV) is sprayed directly onto the punches of the tableting machine. * 0.1% w/w of magnesium stearate is equivalent to 1000 ppm.

Example 3: Properties of the 125 mg dispersible tablet of Example 2

Test	Release specifications
Tablet shape	12 mm diameter, round, flat, bevelled edge with engraving (IA/NVR)
Tablet appearance	Off white
Friability	Max. 1% (0 unit broken)
Crushing strength (mean)	Mean $\geq 70N$
Dissolution rate	Q = 75 % within 30 min
Disintegration time	All units ≤ 3.0 min
Average mass	403.75– 446.25 mg
Mass uniformity	18/20 within $\pm 5\%$ 20/20 within $\pm 10\%$
Content uniformity	10/10 units within 85.0% - 115.0 % / RSD $\leq 6.0\%$
Mean content	95.0 – 105.0 %
Determination of degradation products	unknown degradation product: individual $\leq 0.5\%$ total $\leq 2.0\%$
Fineness of dispersion	Smooth dispersion which passes through a sieve screen with a nominal mesh aperture of 710 μm

Example 4: Properties of the 250 mg dispersible tablet of Example 2

Test	release specifications
Tablet shape	15 mm diameter, round, flat, bevelled edge with engraving (IB/NVR)
Tablet appearance	Off white
Friability	Max. 1% (0 unit broken)
Crushing strength (mean)	Mean $\geq 90N$
Dissolution rate	Q = 75 % within 30 min
Disintegration time	All units < 3 min
Average mass	807.5 – 992.5 mg
Mean content	95.0 – 105.0 %

Example 5: Properties of the 500 mg dispersible tablet of Example 2

Test	Release specifications
Tablet shape	20 mm diameter, round, flat, bevelled edge with engraving (IC/NVR)
Tablet appearance	Off white
Friability	Max. 1% (0 unit broken)
Crushing strength (mean)	Mean \geq 110N
Dissolution rate	Q = 70 % within 30 min
Disintegration time	All units \leq 3.0 min
Average mass	1615 – 1785 mg
Mass uniformity	18/20 within \pm 5% 20/20 within \pm 10%
Mean content	95.0 – 105.0 %
Content uniformity	10/10 units within 85.0% - 115.0 % / RSD \leq 6.0%
Determination of degradation products	unknown degradation product: individual \leq 0.5% total \leq 2.0%
Fineness of dispersion	Smooth dispersion which passes through a sieve screen with a nominal mesh aperture of 710 μ m

Example 6: Magnesium Stearate Assay

	125 mg tablet			250 mg tablet		500 mg tablet		
	L.P.	Pilot	FSP	P.T.	FSP	L.P.	Pilot	FSP
Min (%w/w)	0.1	0.09	0.04	0.08	0.04	0.04	0.03	0.04
Max (%w/w)	0.24	0.36	0.14	0.16	0.08	0.10	0.12	0.06

L.P.: laboratory phase, Pilot: Pilot phase (x 2 of the size of the batch of the laboratory phase), FSP: (batches of production size), P.T.: preliminary trial, 0.1 % w/w of magnesium stearate is equivalent to 1000 ppm.


RSD: relative standard deviation

Example 7: Properties of the 250 mg dispersible tablet of Example 2

Test	Release specifications
Tablet shape	15 mm diameter, round, flat, bevelled edge with engraving (IB/NVR)
Tablet appearance	Off white
Friability	Max. 1% (0 unit broken)
Crushing strength (mean)	Mean $\geq 90N$
Dissolution rate	Q =75 % within 30 min
Disintegration time	All units ≤ 3.0 min
Average mass	807.5 - 892.5 mg
Mass uniformity	18/20 within $\pm 5\%$ 20/20 within $\pm 10\%$
Content uniformity	10/10 units within 85.0% - 115.0 % / RSD $\leq 6.0\%$
Mean content	95.0 -105.0 %
Determination of degradation products	unknown degradation product: individual $\leq 0.5\%$ total $\leq 2.0\%$
Fineness of dispersion	Smooth dispersion which passes through a sieve screen with a nominal mesh aperture of 710 μm

CLAIMS

1. A dispersible tablet comprising Compound I of the formula



or a pharmaceutically acceptable salt thereof present in an amount of from 5% to 40% in weight based on the total weight of the tablet.

2. A dispersible tablet comprising (a) Compound I or a pharmaceutically acceptable salt thereof, and (b) at least one pharmaceutically acceptable excipient suitable for the preparation of tablets, wherein Compound I or a pharmaceutically acceptable salt thereof is present in an amount of from 5% to 40% in weight based on the total weight of the tablet.
3. A dispersible tablet comprising an iron-chelating pharmacologically effective amount of Compound I or a pharmaceutically acceptable salt thereof present in an amount of from 5% to 40% in weight based on the total weight of the tablet.
4. The dispersible tablet according to claim 1, 2 or 3 wherein Compound I is in the free acid form.
5. The dispersible tablet according to any one of claims 1 to 4 wherein Compound I is in a crystalline form.
6. The dispersible tablet according to any one of claims 1 to 5 wherein a lubricant is present in less than 1% in weight based on the total weight of the tablet.
7. The dispersible tablet according to claim 6 wherein the lubricant is present in less than 0.4% in weight based on the total weight of the tablet.

8. The dispersible tablet according to any one of claims 1 to 7 wherein the disintegration time of the tablet is of 5 minutes or less.
9. The dispersible tablet according to any one of claims 1 to 8 wherein the disintegration time of the tablet is of 3 minutes or less.
10. The dispersible tablet according to any one of claims 2 to 9 wherein the pharmaceutically acceptable excipients comprise:
 - (i) at least one filler in a total amount of about 35 to 55 % in weight based on the total weight of the tablet,
 - (ii) at least one disintegrant in a total amount of about 10% to 35% in weight based on the total weight of the tablet
 - (iii) at least one binder in a total amount of about 1.5% to 5% in weight based on the total weight of the tablet,
 - (iv) at least one surfactant in a total amount of about 0.2% to 1% in weight based on the total weight of the tablet,
 - (v) at least one glidant in a total amount of about 0.1% to 0.5% in weight based on the total weight of the tablet, and/or
 - (vi) at least one lubricant in a total amount of less than about 0.4% in weight based on the total weight of the tablet.
11. The dispersible tablet according to any one of claims 6 to 10 wherein the lubricant is magnesium stearate.
12. The dispersible tablet according to anyone of claims 1 to 11 containing Compound I in its free acid form in an amount of about 100 mg to 600 mg .
13. A method of administering to a mammal in need of such a treatment a daily dose of 5 to 40 mg/kg of body weight of Compound I as active ingredient.
14. A process for the preparation of the dispersible tablet according to any one of the preceding claims, which process comprises
 - (i) mixing the Compound I or a pharmaceutically acceptable salt thereof and at least one pharmaceutically acceptable excipient;

- 20 -

- (ii) wet-granulating the mixture obtained in (i);
- (iii) mixing the granulates obtained in (ii) with at least one pharmaceutically acceptable excipient to form a mixture; and
- (iv) spraying the lubricant on the materials contacting surfaces of pressing tools of the tableting machine and compressing the mixture obtained in step (iii) to form a tablet.

15. The process according to claim 14 wherein the lubricant is magnesium stearate.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 03/11351

A. CLASSIFICATION OF SUBJECT MATTER
 IPC 7 A61K9/20 A61K31/41 A61P39/04

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ, BIOSIS, EMBASE, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 97/49395 A (ACKLIN PIERRE ; CIBA GEIGY AG (CH); LATTMANN RENE (CH)) 31 December 1997 (1997-12-31) cited in the application	1-9, 11-15
Y	the whole document page 7, paragraph 2 - page 8, paragraph 2 Examples A-B and D: Tablets comprising resp. 200, 400 and 300 mg of active ingredient	1-15
Y	WO 00/57886 A (TOSELLI DOMINIQUE ; ZAKARIAN NOEL (FR); CLL PHARMA (FR); GIMET RENE (F) 5 October 2000 (2000-10-05) the whole document page 4, lines 12,14,15-21 page 5, lines 1-12 page 6, lines 3-25 page 7, lines 1-26; claims 1,17; examples 1-5	1-15
-/-		

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *Z* document member of the same patent family

Date of the actual completion of the international search

4 March 2004

Date of mailing of the international search report

11/03/2004

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
 NL - 2280 HV Rijswijk
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
 Fax: (+31-70) 340-3016

Authorized officer

Luangkhot, N

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 03/11351

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	& US 2002/061333 A1 (ZAKARIAN ET AL.) 23 May 2002 (2002-05-23) the whole document paragraphs '0018! - '0019!, '0024! - '0025!, '0037! - '0038!, '0042! - '0054!, '0061!; claims 1,17; examples 1-5 -----	1-15
Y	EP 1 070 496 A (KYOWA HAKKO KOGYO KK) 24 January 2001 (2001-01-24) the whole document paragraphs '0001! - '0002!; claims 1,5 -----	14

INTERNATIONAL SEARCH REPORT

International application No.
PCT/EP 03/11351

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

Although claim 13 is directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this International application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 03/11351

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 9749395	A	31-12-1997	AP 1127 A	06-12-2002
			AT 226435 T	15-11-2002
			AU 718037 B2	06-04-2000
			AU 3262997 A	14-01-1998
			BG 103003 A	30-09-1999
			BR 9709973 A	10-08-1999
			CA 2255951 A1	31-12-1997
			CZ 9804272 A3	17-03-1999
			DE 69716602 D1	28-11-2002
			DE 69716602 T2	26-06-2003
			DK 914118 T3	24-02-2003
			WO 9749395 A1	31-12-1997
			EP 0914118 A1	12-05-1999
			ES 2187785 T3	16-06-2003
			HK 1020530 A1	13-06-2003
			JP 2000507601 T	20-06-2000
			KR 2000022222 A	25-04-2000
			NO 986024 A	21-12-1998
			NZ 333308 A	26-05-2000
			PL 330119 A1	26-04-1999
			PT 914118 T	28-02-2003
			RU 2208010 C2	10-07-2003
			SI 914118 T1	30-04-2003
			SK 178598 A3	11-06-1999
			TR 9802693 T2	21-05-1999
			TW 533205 B	21-05-2003
			US 2003069273 A1	10-04-2003
			US 2003203954 A1	30-10-2003
			US 6465504 B1	15-10-2002
WO 0057886	A	05-10-2000	FR 2793690 A1	24-11-2000
			AU 3662700 A	16-10-2000
			BR 0009447 A	08-01-2002
			CA 2368880 A1	05-10-2000
			EP 1165094 A1	02-01-2002
			WO 0057886 A1	05-10-2000
			US 2002061333 A1	23-05-2002
US 2002061333	A1	23-05-2002	FR 2793690 A1	24-11-2000
			AU 3662700 A	16-10-2000
			BR 0009447 A	08-01-2002
			CA 2368880 A1	05-10-2000
			EP 1165094 A1	02-01-2002
			WO 0057886 A1	05-10-2000
EP 1070496	A	24-01-2001	AU 763143 B2	17-07-2003
			AU 5265499 A	01-11-1999
			CA 2327655 A1	21-10-1999
			EP 1070496 A1	24-01-2001
			WO 9952491 A1	21-10-1999